

: 64 1
Vol. 64, No. 1, January, 2003

c-erbB-2 Mitogen-activated Protein Kinases

¹, ²Brain Korea 21 Project, ³

^{1,2} . ² . ² . ¹ . ³ . ⁴

Mitogen-activated Protein Kinases Activities and c-erbB-2 Expression in Breast Cancer Carcinogenesis and Progression

Byeong-Woo Park, MD.^{1,2}, Min-Kyu Heo, B.S.², Ki-Suk Kim, B.S.², Seung-Sang Ko, MD.¹, Seung-Il Kim, MD.³ and Kyong Sik Lee, MD.⁴

Purpose: Increased level mitogen-activated protein kinase (MAPK) and activation of MAPK have been reported in human breast cancers, especially in breast cancers with HER2/neu overexpression. To understand the relationship between the MAPK protein expressions and other clinico-pathological parameters, we examined the status of MAPKs in 20 breast cancers compared to those of paired normals. **Methods:** A total of 20 breast cancers and paired normal breast tissues were included in this study. Tissues were obtained at the operation room and stored at -80°C . Tissue proteins were extracted and the concentration was determined by Bio-Rad protein assay method. Western blot analysis were performed to determine the level of MAPKs expressions using 100 ug of tissue protein in 8%, 10%, or 12% sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE). MAPK assays were carried out by a non-radioactive method developed by Cell Signaling Tech. as recommended by the manufacturer. Clinico-pathological information was provided from the Breast Cancer Registry of Department of Surgery, Yonsei University College of Medicine.

Results: The levels of MAPKs were higher in 95% of breast

cancers compared to those of paired normals. The levels of ERK1/2 were significantly higher in cancer tissues compared to paired normals but the activated forms were not. The levels of JNK, p38, and MKP1 proteins were significantly increased in the cancer tissue compared to the paired normals. The levels of ERK1/2 and activated ERK1/2 proteins were not different between tumor stages. There were no significant differences of the levels of ERK1/2 and activated ERK1/2 proteins between HER2-negative and HER2-positive cancers. There were significantly higher levels of activated ERK1/2 proteins in ER-positive cancers than those in ER-negative cancers ($P < 0.05$).

Conclusion: The levels of MAPKs, but not the activated forms, seem to be increased in breast cancer tissues compared to those of paired normals. The levels of activated MAPKs seem to be associated with estrogen receptor expression in cancer tissues. (*J Korean Surg Soc* 2003;64:6-13)

Key Words: Mitogen-activated protein kinase, HER2/neu, Estrogen receptor, Breast cancer, Stage
: , HER2/neu,
,

Department of ¹Surgery and ²Brain Korea 21 Project, Yonsei University College of Medicine, Department of Surgery, ³Hallym University and ⁴Pochon Chungmun University College of Medicine

: 134
Ⓢ 120-752,
Tel: 02-361-5564, Fax: 02-313-8289
E-mail: bwpark@yumc.yonsei.ac.kr
: 2002 9 17 , : 2002 12 30
1999

가
, 1998
14.1% 가
(I)
가

(apoptosis)) (apoptotic pathway) mi-

c-erbB-2 . togenic signal .(9)

30%

MAPK 가

protein kinases (MAPKs) ,(2,3) Mitogen-actiovated 가

c-erbB-2 MAPK

가 MAPK

(renal cell carcinoma) MAPK

(constitutive activation)가 MAPK

.(4) MAPK mRNA 가 (metastatic

potential)

MAP kinase 가 tyrosyl residue

5 10 가

.(5)

MAP kinase

(dominant oncogenes)

MAP kinase MAP ki-

nase phosphatase (MKP)-1 ras

MAPK-mediated mitogenic effect (6)

.(7) MKP-1

가 , MKP-1

가

ERK-1 MKP-1

가 MKP-1 5q35-ter locus

(loss) MKP-1

(marker)

.(8) MKP-1

MAP kinase

.(8)

MAPK 가 MKP-1

ERK MKP-1 mRNA

JNK

JNK-1

MKP-1 ERK-1 MKP-1

가

MKP-1 JNK

.(9)

1)

1 6 , 2 9 3 5

— 80°C

2)

liquid nitrogen

가 (ice-cold lysis buffer)

(70 mM B-glycerophosphate pH 7.2, 1 mM each mea- and ortho- vanadate, 2 mM manesium chloride, 1 mM EGTA, 1 mM dithiothreitol (DTT), 0.5% Triton X-100, 0.2 mM phenylmethylsulphonyl fluoride (PMSF) and 5 ug ml⁻¹ each of pepstatin A, shymostatin, leupeptin and peptin)

20 (sample) 30

(sonicate) 4°C 23,000 g 15

— 80°C Bradford

(Bio-Rad Laboratories, Richmond, CA, USA)

Bio-Rad

3) Western blot analysis

100 ug

8%, 10%, 12% sodium dodecyl sulphate polyacrylamide gel (SDS-PAGE) Protran Nitrocellulose Membrane (Schleicher and Schuell Corporation, Dassel, Germany) transfer . Blots blocking buffer (20 mM Tris-Cl, pH 7.5, 15 mM sodium chloride, 0.05% Tween-20 (TBST) containing 5% non-fat Carnation milk)

blot (Immunoblot) TBST

1 TBST milk
2 4°C Blots TBST
2 TBST milk 1 2
incubation blots Amersham ECL kit
(Amersham International, Buckinghamshire, UK)
Western band
film GS-690 imaging Densitometer

4) Antibodies

1 Phospho-p44/42 MAP Kinase (Thr202/Tyr204) (Cell Signaling Tech., MA, USA), p44/ 42 MAP Kinase (Cell Signaling Tech., MA, USA), Phospho- p38 MAP Kinase (Thr180/Tyr182) (Cell Signaling Tech., MA, USA), p38 MAP Kinase (Cell Signaling Tech., MA, USA), Phospho-SAPK/JNK (Cell Signaling Tech., MA, USA), SAPK/ JNK (Cell Signaling Tech., MA, USA), MKP-1 (Santa Cruze Biotech., CA USA) p44/42 MAP Kinase Assay Kit (Cell Signaling Tech., MA, USA) In vitro MAPK assay
2 Anti-mouse IgG-HRP (Santa Cruze Biotech., CA USA) Anti-rabbit IgG-HRP

(Santa Cruze Biotech., CA USA)

5) In Vitro MAPK assays

MAPK assays Cell Signaling Tech. (MA, USA) 71
MAPKs 300
ug phospho-specific ERK 1/2 MAPK (Thr202/ Tyr204) (immunoprecipitation)
kinase assays ATP substrate Elk-1
phospho-Elk-1 phospho-specific Elk-1 (Ser 383) Western blot analysis
phospho Elk-1 Ser383 Elk-1

6)

SPSS 10.0
SPSS
independent sample t-test
crosstab P < 0.05




Fig. 1. Levels of MAPKs in the representative breast cancers and paired normals. The ERK 1/2 proteins were detected by Western blot analysis by using ERK 1/2 polyclonal antibody (upper panel). The phospho-ERK 1/2 proteins were detected by Western blot analysis by using phospho-specific MAPK (pERK 1/2) antibody (lower panel). The levels of ERK 1/2 were significantly higher in cancer tissues compared to paired normals but the activated forms were not. N and C represent normal and carcinoma, respectively.




Fig. 2. Levels of JNK, p38, and MKP1 in the representative breast cancers and paired normals. Each protein was detected by Western blot analysis by using JNK 1/2 polyclonal antibody (uppermost panel), p38 polyclonal antibody (middle panel), and MKP1 polyclonal antibody (lowest panel). The levels of JNK, p38, and MKP1 proteins were significantly increased in the cancer tissue compared to the paired normals. N and C represent normal and carcinoma, respectively.

1) MAPKs proteins and activated ERK1/2 in cancer tissues and paired normals

ERK1/2
(5%)
12.2)
ERK1/2
ERK1/2 6 (30%)
8 (40%) 가 6 (30%) 가
1.17
(0.21 5.6) 가
MAPK
MAPK
JNK
가 , p38
MAPK
1.71 (0.5 3.26) 가 MAPK
가 (Table 1, Fig. 1, 2).

2) Levels of ERK1/2 and activated ERK1/2 according to the axillary lymph node status and tumor stages

MAPKs 3 ERK1/2 가 (Table 1, Fig. 3)

3) Levels of ERK1/2 and activated ERK1/2 according to HER2 and ER expressions

HER2 MAPK
(Table 1, Fig. 4), ER ERK1/2
가 ERK1/2 (P < 0.05) 가 (Table 1, Fig. 5).

4) Levels of ERK1/2 and activated ERK1/2 according to histologic grade and MKP1 levels

ERK1/2

Table 1. Clinico-pathological characteristics and the ratios of MAPKs in cancer tissues over paired normals

Sample	LN	Stage	HG	ER	HER2	ERK	pERK	JNK	p38	MKP1
1	0	1	2	0	1	1.1	0.8		1.5	2.7
2	1	2	2	1	0	1.9	0.9		1.8	2.4
3	0	1		1	0	1.8	1.5		1.0	0.5
4	2	2		1	1	2.6	1.4	3.1	2.3	1.8
5	0	2	3			6.6	1.0	1.5	9.6	1.9
6	1	2	2	1	0	7.0	2.5	2.4	8.4	1.2
7	2	2	2	1	0	2.5	1.2	1.7	3.5	2.1
8	0	2	2	1	1	12.2	1.0	2.3	5.5	1.8
9	0	2	2	0	0	4.3	0.9	1.4	1.3	0.8
10	26	2	2	0	1	3.6	0.9	2.9	2.1	
11	0	1	3	0	0	4.2	0.6	2.0	14.6	3.3
12	1	2	1	1	0	1.1	0.8	1.2	1.0	
13	18	3	3	0	1	1.0	0.5	1.2	1.3	1.0
14	15	3	3	0	1	2.2	0.2	1.0	1.7	1.8
15	0	2	2	1	1	11.1	5.6	3.3	4.1	0.7
16	0	1	3	0	0	9.1	0.4	1.7	3.1	1.2
17	11	3	2	1	0	0.7	1.5	0.8	1.0	
18	0	1	2	0	1	3.7	0.3	2.0	25.3	2.0
19	4	3	3	1	0	6.8	1.0	2.2	9.0	2.0
20	0	1	2	1	1	2.5	0.4	1.3	3.0	1.8

LN, HG, and ER represents the number of metastatic lymph nodes in the axilla, histologic grade, and estrogen receptor, respectively. The values of MAPKs were calculated by denomination of the optical density of Western blots of Cancer tissue over that of paired normal tissue.




Fig. 3. Levels of MAPKs in the representative breast cancers and paired normals according to tumor stages. The ERK 1/2 proteins were detected by Western blot analysis by using ERK 1/2 polyclonal antibody (left panel). The phospho-ERK 1/2 proteins were detected by Western blot analysis by using phospho-specific MAPK (pERK 1/2) antibody (right panel). Uppermost panel represents stage 1, middle one represents stage 2, and lowest panel represents stage 3. The levels of ERK 1/2 and activated ERK 1/2 proteins were not different between stages. N and C represent normal and carcinoma, respectively.




Fig. 4. Levels of MAPKs in the representative breast cancers and paired normals according to HER2 expressions. The ERK 1/2 proteins were detected by Western blot analysis by using ERK 1/2 polyclonal antibody (upper panel). The phospho-ERK 1/2 proteins were detected by Western blot analysis by using phospho-specific MAPK (pERK 1/2) antibody (lower panel). There were no significant differences of the levels of ERK 1/2 and activated ERK 1/2 proteins between HER2-negative (left panel) and HER2-positive cancers. N and C represent normal and carcinoma, respectively.




Fig. 5. Levels of MAPKs in the representative breast cancers and paired normals according to ER expressions. The ERK 1/2 proteins were detected by Western blot analysis by using ERK 1/2 polyclonal antibody (upper panel). The phospho-ERK 1/2 proteins were detected by Western blot analysis by using phospho-specific MAPK (pERK 1/2) antibody (lower panel). There were significantly higher levels of activated ERK 1/2 proteins ER-positive (left panel) cancers than those in ER-negative cancers ($P < 0.05$). N and C represent normal and carcinoma, respectively.

(Table 1) MKP1

가
(60% vs 33%) ERK

HER2

ERK 1/2
1, Fig. 6).

가

($P=0.086$)

(Table




Fig. 6. Levels of MAPKs in the representative breast cancers and paired normals according to MKP1 expression. There were no significant differences in ERK1/2 and activated forms between higher MKP1 (samples 4, 5, and 8) and Lower MKP1 groups (sample 9). N and C represent normal and carcinoma, respectively.

MAPK (20-23) MAPK

가 ERK1/2

MAPK (fibroblast) MAPK kinase (MEK) 가 HER2

MAPK oncogenicity ER (P < 0.05),

MAPK (10) MAPK (activated form) MAPK 가 가 ER

MAPK HER2 MAPK

MAPK MAPK 가

MAPK , MAPK ER MAPK ; ,

MAPK 가 ERK JNK p38 (11) MAPK 가; ,

knase activated form MAPK (19) (24-26)

MAPK Mueller(11) MAPK

MAPK MAPK 가 ERK (11) 가 MAPK

MAPK MAPK 가

MAPK (12-15) ERK JNK

MAPK (hydrogen peroxide) p38

ERK1/2 (16) ER ERK1/2 가 가

HER2 (major regulator) (17) HER1 p38 JNK

가 (18) ras MAPK

aggressive tumor phenotype (14) (an ERK1/2 JNK

MAPK 가

(19) MAPK 가 ERK1/2

(18) , potent mi- ERK1/2

togenic stimuli 가 (4.31 : 20

19 1.01 12.21 가) ERK 1/2 ERK 1/2
ER (P < 0.05) ERK 1/2
MKP1 MKP1 가
MKP1 MAPK 가
(8)
MKP1 가 ERK 1/2
(P=0.086)
가
stress-activated protein kinase p38 JNK
가
MAPK
MAPK
mapk
5 10 MAPK 가 .(5)
mapk 5 20 가
(5) MAPK
mRNA (metastatic potential)
가 .(27) MAPK
(4) ERK 1/2
가
MAPK
MKP1 MAPK MAPK
mitogenic signal
MAPK mitogenic effect
(6) (block differentiation).(7) MKP1
mitogenic signals, MAPK
(28) MKP1
가
MKP1
MKP1
ERK 1
MKP1
MKP1
(8) MKP1 mRNA
가 ERK 1, 2 가
MKP1
(high-grade
tumor) MKP1 ERKs
MAPK (MAPK
independence) .(29,30)
MAPK
(8) MKP1 HER2 , ER
, ERK 1/2

REFERENCES

- 1) The Korean Breast Cancer Society. Clinical characteristics of Korean breast cancer patients in 1998. *J Korean Med Sci* 2000; 15:569-79.
- 2) Slamon DJ, Clark GM, Wong SG, Levin WJ, Ullrich A, McGuire WL. Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene. *Science* 1987;235:177-82.
- 3) Slamon DJ, Godolphin W, Jones LA, Holt JA, Wong SG, Keith DE, et al. Studies of the HER-2/neu proto-oncogene in human breast and ovarian cancer. *Science* 1989;244:707-12.
- 4) Oka H, Chatani Y, Hoshino R, Ogawa O, Kakehi Y, Terachi T, et al. Constitutive activation of mitogen-activated protein (MAP) kinases in human renal cell carcinoma. *Cancer Res* 1995;55(18):4182-7.
- 5) Sivaraman VS, Wang H, Nuovo GJ, Malbon CC. Hyperexpression of mitogen-activated protein kinase in human breast cancer. *J Clin Invest* 1997;99:1478-83.
- 6) Sun H, Tonks NK, Bar-Sagi D. Inhibition of Ras-induced DNA synthesis by expression of the phosphatase MKP-1. *Science* 1994;266:285-8.
- 7) Yao H, Labudda K, Rim C, Capodiceci P, Loda M, Stork PJ. Cyclic adenosine monophosphate can convert epidermal growth factor into a differentiating factor in neuronal cells. *J Biol Chem* 1995;270:20748-53.
- 8) Loda M, Capodiceci P, Mishra R, Yao H, Corless C, Grigioni W, et al. Expression of mitogen-activated protein kinase phosphatase-1 in the early phases of human epithelial carcinogenesis. *Am J Pathol* 1996;149:1553-64.
- 9) Magi-Galluzzi C, Mishra R, Fiorentino M, Montironi R, Yao H, Capodiceci P, et al. Mitogen-activated protein kinase phosphatase 1 is overexpressed in prostate cancers and is inversely related to apoptosis. *Lab Invest* 1997;76:37-51.
- 10) Brunet A, Pages G, Pouyssegur J. Constitutively active mutants of MAP kinase kinase (MEK1) induce growth factor-relaxation and oncogenicity when expressed in fibroblasts. *Oncogene* 1994;9:3379-87.
- 11) Mueller H, Flury N, Eppenberger-Castori S, Kueng W, David F, Eppenberger U. Potential prognostic value of mitogen-activated protein kinase activity for disease-free survival of primary breast cancer patients. *Int J Cancer* 2000;89:384-8.
- 12) Blenis J. Signal transduction via the MAP kinases: proceed at

- your own RSK. *Proc Natl Acad Sci USA* 1993;90:5889-92.
- 13) Cobb MH, Goldsmith EJ. How MAP kinases are regulated. *J Biol Chem* 1995;270:14843-6.
- 14) Janes PW, Daly RJ, deFazio A, Sutherland RL. Activation of the Ras signalling pathway in human breast cancer cells over-expressing erbB-2. *Oncogene* 1994;9:3601-8.
- 15) Callans LS, Naama H, Khandelwal M, Plotkin R, Jardines L. Raf-1 protein expression in human breast cancer cells. *Ann Surg Oncol* 1995;2:38-42.
- 16) Stevenson MA, Pollock SS, Coleman CN, Calderwood SK. X-irradiation, phorbol esters, and H₂O₂ stimulate mitogen-activated protein kinase activity in NIH-3T3 cells through the formation of reactive oxygen intermediates. *Cancer Res* 1994; 54:12-5.
- 17) Pearson G, Robinson F, Beers Gibson T, Xu BE, Karandikar M, Berman K, et al. Mitogen-activated protein (MAP) kinase pathways: regulation and physiological functions. *Endocr Rev* 2001;22:153-83.
- 18) von Lintig FC, Dreilinger AD, Varki NM, Wallace AM, Casteel DE, Boss GR. Ras activation in human breast cancer. *Breast Cancer Res Treat* 2000;62:51-62.
- 19) Santen RJ, Song RX, McPherson R, Kumar R, Adam L, Jeng MH, et al. The role of mitogen-activated protein (MAP) kinase in breast cancer. *J Steroid Biochem Mol Biol* 2002;80:239-56.
- 20) Adams JM, Cory S. The Bcl-2 protein family: arbiters of cell survival. *Science* 1998;281:1322-6.
- 21) Datta SR, Dudek H, Tao X, Masters S, Fu H, Gotoh Y, Greenberg ME. Akt phosphorylation of BAD couples survival signals to the cell-intrinsic death machinery. *Cell* 1997;91: 231-41.
- 22) Kauffmann-Zeh A, Rodriguez-Viciana P, Ulrich E, Gilbert C, Coffey P, Downward J, et al. Suppression of c-Myc-induced apoptosis by Ras signalling through PI(3)K and PKB. *Nature* 1997;385:544-8.
- 23) Nass SJ, Dickson RB. Defining a role for c-Myc in breast tumorigenesis. *Breast Cancer Res Treat* 1997;44:1-22.
- 24) Cicatiello L, Addeo R, Altucci L, Belsito Petrizzi V, Boccia V, Cancemi M, et al. The antiestrogen ICI 182,780 inhibits proliferation of human breast cancer cells by interfering with multiple, sequential estrogen-regulated processes required for cell cycle completion. *Mol Cell Endocrinol* 2000;165:199-209.
- 25) Addeo R, Altucci L, Battista T, Bonapace IM, Cancemi M, Cicatiello L, et al. Stimulation of human breast cancer MCF-7 cells with estrogen prevents cell cycle arrest by HMG-CoA reductase inhibitors. *Biochem Biophys Res Commun* 1996; 220:864-70.
- 26) Altucci L, Addeo R, Cicatiello L, Dauvois S, Parker MG, Truss M, et al. 17beta-Estradiol induces cyclin D1 gene transcription, p36D1-p34cdk4 complex activation and p105Rb phosphorylation during mitogenic stimulation of G(1)-arrested human breast cancer cells. *Oncogene* 1996;12:2315-24.
- 27) Pencil SD, Toh Y, Nicolson GL. Candidate metastasis-associated genes of the rat 13762NF mammary adenocarcinoma. *Breast Cancer Res Treat* 1993;25:165-74.
- 28) Sun H, Charles CH, Lau LF, Tonks NK. MKP-1 (3CH134), an immediate early gene product, is a dual specificity phosphatase that dephosphorylates MAP kinase in vivo. *Cell* 1993; 75:487-93.
- 29) Fearon ER, Vogelstein B. A genetic model for colorectal tumorigenesis. *Cell* 1990;61:759-67.
- 30) Kastirnakis WV, Ramchurren N, Rieger KM, Hess DT, Loda M, Steele G, et al. Increased incidence of p53 mutations is associated with hepatic metastasis in colorectal neoplastic progression. *Oncogene* 1995;11:647-52.